

COMPLEX CHRONIC DISEASES PROGRAM Medication Handout

Duloxetine (Cymbalta)

Date: May 15, 2018

Medication: Duloxetine (Cymbalta) 30 mg

What is Duloxetine:

Duloxetine is a type of medication called "Serotonin Norepinephrine Reuptake Inhibitors" or "SNRIs" that was initially developed to treat depression. It works by altering the levels of certain transmitters in the brain such as noradrenalin and serotonin.

Duloxetine can also be used in the treatment of anxiety, irritable bowel syndrome (with constipation), pain, and certain symptoms of myalgic encephalomyelitis / chronic fatigue syndrome (but not the fatigue itself).

Expected Benefit:

• Usually takes 4 – 6 weeks to notice a benefit

What to expect:

- "Transition" effects
 - These symptoms are the effects of your body getting used to the medication (i.e., transition onto the medication)
 - It's different for everybody. Some people feel "not like themselves," a bit anxious, or unusual.
 - These symptoms typically go away after 7 to 10 days
 - o Please call the CCDP if the anxiety is intolerable.

Watch for possible side effects:

This list of side effects is important for you to be aware of however, it is also important to remember that not all side effects happen to everyone.

If you have problems with these side effects talk with your doctor or pharmacist:

- Trouble sleeping that is more than usual
- Heart burn, stomach upset
- Sexual dysfunction
- High blood pressure (Make sure you family doctor is regularly checking your blood pressure)

Stopping the medication:

This medication should not be stopped abruptly. Your doctor may advise you to reduce the dose slowly to help prevent your symptoms from returning.

How to use this medication:

Take this medication with or without food

Dosing Schedule:

- Start with 30 mg every 2nd day
- Increase dose according to table below



COMPLEX CHRONIC DISEASES PROGRAM Medication Handout

Duloxetine (Cymbalta)

Date: May 15, 2018

You can stay at the same dose (stop increasing) if you get side effects

0	You	might	want to	lower	the	dose	one	step
---	-----	-------	---------	-------	-----	------	-----	------

AM				
30 mg every 2 nd day	For 2 weeks			
30 mg daily	For 1 month; stay on this dose			
Follow up with clinic before increasing the dose				

Drugs and Foods to Avoid:

Ask your doctor or pharmacist before using any other medication, including non-prescription medication (over-the-counter medication) and herbal products.

- Avoid dextromethorphan ("DM" cough products)
- Avoid alcohol use
- Avoid chamomile, primrose oil, fenugreek, flaxseed, 5 HTP, feverfew, fish oils, omega 3, flaxseed, ginkgo, ginseng, red clover, SAMe, willow bark, glucosamine, green tea, horse chestnut seed, St John's Wort

Please contact the Complex Chronic Diseases Program at if you have further questions about your medications.