

### COMPLEX CHRONIC DISEASES PROGRAM

### Medication Handout

Date: May 15, 2018

### Medication: Zolmitriptan (Zomig) 2.5 mg or 5 mg

#### What is Zolmitriptan:

Zolmitriptan is a type of medication called “triptans” which is used to treat acute migraine headaches. Migraine symptoms may be caused by swollen blood vessels in the head. Zolmitriptan is thought to reduce the swelling of these blood vessels. This helps to take away the headache and other symptoms of a migraine attack, such as feeling or being sick (nausea or vomiting) and being sensitive to light and sound.

Zolmitriptan works only when a migraine attack has started. It will not stop you from getting an attack

#### Expected Benefit:

- Relief should occur within 1 – 2 hrs

#### Watch for possible side effects:

This list of side effects is important for you to be aware of however, it is also important to remember that not all side effects happen to all people.

If you have problems with these side effects talk with your doctor or pharmacist

- Heaviness of the arms and legs
- Flushing
- Sensation of pins and needles
- Tightness in the chest, neck, or throat
- These side effects can be reduced or avoided with a smaller dose or trying a different triptan (e.g., rizatriptan)
- If you develop muscle spasms and fever, stop the medication and let the clinic know

#### Stopping the medication:

There should be no withdrawal effect when stopping the medication as the medication is not taken regularly.

#### How to use and store this medication:

- Take this medication with or without food
- Most effective if taken at the earliest onset of migraine
- Can be taken with NSAIDs (e.g., ibuprofen, naproxen, diclofenac)
- Can also be taken with metoclopramide if nausea is severe

#### Dosing Schedule:

- 2.5 mg or 5 mg at earliest onset of headache
- Dose may be repeated in 2 hrs if no benefit
- Maximum 10 mg per day
- Should not be taken on a daily basis

Date: May 15, 2018

- Use of analgesics for migraines more than 10 times per month increases the risk of “Medication Overuse Headaches”
- Frequent migraines may benefit from medication for migraine prevention

**Drug and Foods to Avoid:**

Ask your doctor or pharmacist before using any other medication, including non-prescription medication (over-the-counter medication) and herbal products.

- Do not take with St. John’s wort

Please contact the Complex Chronic Diseases Program if you have further questions about your medications.